arXiv:1403.4381v2 [math.CT] 31 Mar 2014

PROPERNESS AND SIMPLICIAL RESOLUTIONS FOR THE MODEL CATEGORY dgCat

JULIAN V. S. HOLSTEIN

1. Introduction

We provide proofs of the following properties of the model category **dgCat** of dg-categories (with the Morita or Dwyer-Kan model structure).

- When k is a field, the category $\mathbf{dgCat}_k$ is left proper.
- Natural simplicial resolutions in **dgCat** are given by dg-categories of Maurer-Cartan elements.

Left properness is essential to show the existence of Bousfield localizations of dg-categories. We also remark that **dgCat** is cellular and there is a Quillen equivalent combinatorial subcategory (without assumptions on the existence of large cardinals).
Fernando Muro has pointed out that left properness already follows from his more general results, namely Corollary 1.3 in [7].
Simplicial resolutions allow for constructions of explicit mapping spaces and simplicial actions. These play a crucial role in categorifying cohomology to Morita cohomology, see [4]. We construct simplicial resolutions by an explicit if somewhat lengthy computation motivated by the Čech globalization in [10]. Note that the explicit combinatorics of this construction have appeared in other contexts: If K is the nerve of a category this is the data of an A_∞-functor, see for example [5]. If K is any simplicial set one recovers the ∞-local systems defined in [1]. We feel that the interpretation here as the cotensor action of simplicial sets on **dgCat**, computed via simplicial resolutions, provides a satisfying conceptual viewpoint.
These results are taken from the author's thesis. Thanks are due to Ian Grojnowski and Jon Pridham for helpful discussions.

1.1. Conventions.

We assume the reader is familiar with the theory of dg-categories. Basic references are [6] and [12].
Recall in particular that there are two model structures on $\mathbf{dgCat}_k$, the category of differential graded categories over a ring k. These are the Dwyer-Kan model structure and its left Bousfield localization, the Morita

model structure. We will often not distinguish between them as our results will apply to both model categories.
We use homological grading conventions, all differentials decrease the degree. The degree is indicated by a subscript or the inverse of a superscript, $C_i = C^{-i}$.

2. Dg-categories over a field form a left proper, cellular, combinatorial model category

2.1. **Left properness.** In this section we will show that the model category of dg-categories over a field k is left proper. Recall that a model category is *left proper* if any pushout of a weak equivalence along a cofibration is again a weak equivalence.

Remark 1. Recall that **dgCat** with the Dwyer-Kan model structure is right proper since every object is fibrant, and it is not right proper with the Morita model structure, as is shown explicitly by Example 4.10 in [11].

Before proceeding to the proof we mention that Dwyer and Kan prove left properness for simplicial categories on a fixed set of objects in [2].

Theorem 1. *If k is a field the model category* $\mathbf{dgCat}_k$ *is left proper.*

Proof. Left Bousfield localization preserves left properness, see Proposition 3.4.4 of [3], so it is enough to show **dgCat** with the Dwyer-Kan model structure is left proper.
The main work is in showing that pushout along the generating cofibrations preserves quasi-equivalences.
To see this suffices note first that transfinite compositions are just filtered colimits, and filtered colimits preserve quasi-equivalences as follows: A filtered colimit of categories can be computed set-theoretically on objects and morphisms. Now filtered colimits preserve weak equivalences of simplicial sets and hence of mapping spaces. They also preserve the homotopy category since a filtered colimit of equivalences of categories is an equivalence of categories and taking the homotopy category commutes with filtered colimits. Second, if pushout along some map preserves weak equivalences then so does pushout along a retract by functoriality of colimits. Since all cofibrations are retracts of transfinite compositions of generating cofibrations, it does indeed suffice to check generating cofibrations.
Recall the generating cofibrations of **dgCat** [11]. We write k for the dg-catgeory with one object with endomorphisms k concentrated in degree 0. Also let $\mathscr{S}(n)$ have two objects a and b and $\mathrm{End}(a) \cong \mathrm{End}(b) \cong k[0]$ while $\mathrm{Hom}(a,b) = k.g$ with g in degree n and $\mathrm{Hom}(b,a) = 0$. Finally let $\mathscr{D}(n)$ be obtained by $\mathscr{S}(n)$ by adding a generating morphism f of degree $n+1$ to

$\mathscr{S}(n)$ with $df = g$. Then the generating cofibrations of **dgCat** are given by $\emptyset \to k$ and by $\mathscr{S}(n) \to \mathscr{D}(n)$ for all n.
It is clear that pushout along $\emptyset \to k$ preserves quasi-equivalences.
So consider the generating cofibration $\mathscr{S}(n) \to \mathscr{D}(n)$ with a map $j\colon \mathscr{S}(n) \to \mathscr{C}$ and a quasi-equivalence $F\colon \mathscr{C} \to \mathscr{E}$. In forming the pushforward we adjoin a new map f with $df = j(g)$. We call the resulting category $\mathscr{C}'$. Then let $\mathscr{E}'$ be the pushout of $\mathscr{S}(n) \to \mathscr{D}(n)$ along $F \circ j$.
The pushout along j has the same objects as $\mathscr{C}$. The morphism space is obtained by collecting maps from C to D, graded by how often they factor through $f\colon j(a) \to j(b)$. Write $\mathscr{C}(A, B)$ etc. for the enriched hom-spaces $\underline{\mathrm{Hom}}_{\mathscr{C}}(A, B)$ etc. Then the hom-spaces in $\mathscr{C}'$ are given as follows:

$$\mathscr{C}'(C, D) = \mathrm{Tot}^{\oplus}\left(\mathscr{C}(C, D) \oplus (\mathscr{C}(j(b), D) \otimes k.f \otimes T \otimes \mathscr{C}(C, j(a)))\right) \tag{1}$$

Here $T = \sum_{n\geq 0}(\mathscr{C}(j(b), j(a)) \otimes k.f)^{\otimes n}$ and we introduce a horizontal degree n with $\mathscr{C}(C, D)$ in degree -1. The right hand side has a vertical differential given by the internal differential and a horizontal differential given by $f \mapsto j(g) \in \mathrm{Hom}(j(b), j(a))$ composed with the necessary compositions.
If the functor F is not the identity on objects from $\mathscr{C}$ to $\mathscr{E}$ we factor

$$F = Q \circ H\colon \mathscr{C} \to \mathscr{D} \to \mathscr{E}$$

where $\mathscr{D}$ has as objects the objects of $\mathscr{C}$ but $\mathrm{Hom}_{\mathscr{D}}(A, B) = \mathrm{Hom}_{\mathscr{E}}(FA, FB)$. Then H is identity on objects and Q is an isomorphism on hom-spaces. We form the pushforward and obtain the factorization $F' = Q' \circ H'$ through $\mathscr{D}'$. So it suffices to prove the following two lemmas. □

Lemma 2. *The functor Q' defined as above is a quasi-equivalence if Q is.*

Proof. Note that Q' is quasi-essentially surjective if Q is since both $\mathscr{D} \to \mathscr{D}'$ and $\mathscr{E} \to \mathscr{E}'$ are essentially surjective as pushout along j does not change the set of objects.
We filter the right-hand side of Equation 1 for hom-spaces in $\mathscr{E}'$ by columns, i.e. by n. This filtration is bounded below and exhaustive for the direct sum total complex and hence the spectral sequence converges. Now we use the fact that Q induces isomorphisms on hom-spaces to obtain an isomorphism of spectral sequences, which in turn induces an isomorphism $\mathscr{D}'(C, D) \cong \mathscr{E}'(QC, QD)$, so Q' gives weak equivalences on mapping spaces.
Note that since $\mathscr{S}(n)$ maps to $\mathscr{E}$ via $\mathscr{D}$ all the hom-spaces involved in computing $\mathscr{E}'(QC, QD)$ are indeed images of hom-spaces in $\mathscr{D}$. □

Lemma 3. *The functor H' defined as above is a quasi-equivalence if H is.*

Proof. Note that H' is quasi-essentially surjective if H is for the same reason that Q' is.
To consider the effect of H' on mapping spaces proceed as in the previous lemma. Now H only induces weak equivalences on hom-spaces, but we

know all hom-spaces are flat over k as k is a field. Hence the tensor product in Equation 1 preserves weak equivalences. So the same spectral sequence argument applies. □

Remark 2. If we have some non-flat hom-spaces then the conclusion is false. Consider Example 2.11 in [9] for the case of dg-algebras. In that paper the existence of a proper model for simplicial k-algebras is proven. A similar result for dg-categories is beyond the scope of this work.
Hence the model category of dg-categories is only left proper if all dg-categories are k-flat, i.e. if and only if k has flat dimension 0, which is the case if and only if k is von Neumann regular.

2.2. **Cellularity and combinatoriality.** One of the main uses of properness is in constructing left Bousfield localizations. The only additional assumption needed is that the model category is either cellular or combinatorial. We now show that both are satisfied for **dgCat**.

Proposition 4. *The two model category structures on* **dgCat** *are cellular.*

Proof. Recall that a model category is cellular if it is cofibrantly generated with generating cofibrations I and generating trivial cofibrations J such that the domains and codomains of the elements of I are compact, the domains of the elements of J are small relative to I and the cofibrations are effective monomorphisms. See Chapter 10 of [3] for more details.
Left Bousfield localization preserves being cellular see Theorem 4.1.1 of [3]. So it is enough to show **dgCat** with the Dwyer-Kan model structure is cellular.
The domains and codomains of elements of I are categories with at most two objects and perfect hom-spaces, so maps from these objects to relative I-complexes factor through small subcomplexes. So domains and codomains of I are compact.
Similarly the domains of the elements of J have two objects and perfect hom-spaces. Hence taking maps from a domain of J commutes with filtered colimits. So domains of J are small relative to I.
We are left to check that relative I-cell complexes, i.e. transfinite compositions of pushouts of generating cofibrations, are effective monomorphisms, i.e. any relative I-cell complex $f\colon X \to Y$ is the equalizer of $Y \rightrightarrows Y \amalg_X Y$.
Note that we form the pushout along a generating cofibration by attaching maps freely. If we form $\mathscr{C}'$ and $\mathscr{C}''$ from $\mathscr{C}$ by attaching maps freely then the equalizer will have the same objects and the hom-spaces are given by considering morphisms of the pushout that are in the image of both $\mathscr{C}'$ and $\mathscr{C}''$. But these are precisely the hom-spaces of $\mathscr{C}$. □

Definition. Let λ be a regular cardinal. An object A in a category $\mathscr{D}$ is *λ-presentable* if it is small with respect to λ-filtered colimits, i.e. if for every

λ-filtered colimit $\operatorname{colim} B_i$ the map $\operatorname{colim} \operatorname{Hom}(A, B_i) \to \operatorname{Hom}(A, \operatorname{colim} B_i)$ is an isomorphism. We say A is *presentable* if it is λ-presentable for some λ. A cocomplete category is *locally presentable* if for some regular cardinal λ it has a set S of λ-presentable objects such that every object is a λ-directed colimit of objects in S.

Definition. A model category is *combinatorial* if the underlying category is locally presentable.

It is known that there exist combinatorial models for all cofibrantly generated model categories under a large cardinal assumption, cf. [8]. We notice that this assumption is not necessary for **dgCat**.

Proposition 5. *The category* **dgCat** *is Quillen equivalent to a combinatorial subcategory.*

Proof. This follows immediately from the proof of the main theorem of [8]. Let $\mathscr{D}$ denote either of the two model structures on **dgCat**. Let S be the collection of objects that are domains or codomains of the generating cofibrations and generating trivial cofibrations. (See [11] for an explicit description.) Clearly S is a set. Let $\mathscr{S}$ denote the full subcategory of $\mathscr{D}$ with objects S. Define $\eta_S(X)$ to be the colimit of the forgetful diagram $(s \to A) \mapsto s$ indexed by the overcategory $\mathscr{S} \downarrow A$. Then an object $A \in \mathscr{D}$ is *S-generated* if it is isomorphic to $\eta_S(X)$.
Now by the proof of Theorem 1.1 in [8] the subcategory of S-generated objects of $\mathscr{D}$ is a model category $\mathscr{D}_S$ which is Quillen equivalent to the original one. Moreover, by Proposition 3.1 of [8], $\mathscr{D}_S$ is locally presentable if every object in S is presentable. But this is clear since the objects in S have finitely many objects and generating morphisms. □

Remark 3. Note that Vopěnka's principle is not needed here since the objects of S are presentable.

3. Simplicial resolutions of dg-categories

In this section we will construct explicit simplicial functorial resolutions $\mathscr{C} \mapsto \mathscr{C}_\bullet$ in **dgCat**. Again, we can consider either model structure on **dgCat**.
We first recall the basic definitions. Let Δ be the simplex category and consider the constant diagram functor $c\colon \mathscr{M} \to \mathscr{M}^{\Delta^{\mathrm{op}}}$. Then a *simplicial resolution* $M_\bullet$ for $M \in \mathscr{M}$ is a fibrant replacement for cM in the Reedy model structure on $\mathscr{M}^{\Delta^{\mathrm{op}}}$. (For a definition of the Reedy model structure see for example Chapter 15 of [3].) The dual notion is a *cosimplicial resolution* $M^\bullet$.
We recall two applications:

By using simplicial resolutions one can define mapping spaces with values in $Ho(\mathbf{sSet})$ for every model category, even if it is not a simplicial model category. If $cB \to \tilde{B}$ is a simplicial resolution in $\mathscr{M}^{\Delta^{\mathrm{op}}}$ and QA a cofibrant replacement in $\mathscr{M}$ then $\mathrm{Map}(A,B) \simeq \mathrm{Hom}^\bullet(QA,\tilde{B}) \simeq R(\mathrm{Hom}^\bullet(-,c-))$, where the right-hand side uses the bifunctor $\mathrm{Hom}^\bullet \colon \mathscr{M}^{\mathrm{op}} \times \mathscr{M}^{\Delta^{\mathrm{op}}} \to \mathbf{Set}^{\Delta^{\mathrm{op}}}$ that is defined levelwise.
Moreover, every homotopy category of a model category is tensored and cotensored in $Ho(\mathbf{sSet})$. In fact, $\mathscr{M}$ can be turned into a simplicial category in the sense that there is an enrichment given by the bifunctor Map and there is a tensor functor as well as a cotensor or power functor, which can be constructed from the simplicial and cosimplicial resolutions. The cotensor is constructed using the simplicial resolution as follows: Let a simplicial resolution $A_\bullet \in \mathscr{M}^{\Delta^{\mathrm{op}}}$ and a simplicial set K be given. Consider ΔK^{op}, the opposite of the category of simplices of K, with the natural map $v \colon \Delta K^{\mathrm{op}} \to \Delta^{\mathrm{op}}$ sending $\Delta[n] \mapsto K$ to $[n]$. We define $A_\bullet^K = \lim_{\Delta K^{\mathrm{op}}} A_n$ to be the image of $A_\bullet$ under $\lim \circ\, v^* \colon \mathscr{C}^{\Delta^{\mathrm{op}}} \to \mathscr{C}^{\Delta K^{\mathrm{op}}} \to \mathscr{C}$. This can also be written as $A^K = \lim_n (\prod_{K_n} A_n)$.

3.1. **The construction.** Our construction is directly motivated by Simpson's construction of the globalization of a presheaf of dg-categories as a dg-category of Maurer–Cartan elements, cf. section 5.4 of [10].

Remark 4. In fact, the construction of $\mathscr{C}_n$ below corresponds to considering the constant presheaf of dg-categories on a covering of $|\Delta^n|$ by $n+1$ open sets (corresponding to leaving out one of the faces).

Definition. Assume $\mathscr{C}$ is fibrant, replace fibrantly otherwise. Then $\mathscr{C}_n$ is a dg-category with objects given by pairs (E,η) where E is a collection $E_0,\dots,E_n \in \mathrm{Ob}\mathscr{C}$ and η is a collection of $\eta_I = \eta(I) \in \underline{\mathrm{Hom}}_{k-1}(E_{i_0},E_{i_k})$ for all multi-indices $I = (i_0,\dots,i_k)$ with $1 \le k \le n$. The case $k=0$ is subsumed by the differential on E. (We interpret $\eta(i) = 0$ where it comes up in computation.) These pairs must satisfy the Maurer–Cartan condition: $\delta\eta + \eta^2 = 0$, explained below. We also demand that all $\eta_i \in \underline{\mathrm{Hom}}(E_i,E_i)$ are weak equivalences in $\mathscr{C}$.

Remark 5. If we do not fibrantly replace the construction gives a *simplicial framing* on **dgCat**, see for example [3]. The simplicial resolution can then be viewed as composing functorial fibrant replacement with the simplicial framing.

Let us spell out the Maurer–Cartan condition. Intuitively, η provides all the comparison maps as well as homotopies between the different

compositions. We define the differential

$$(\delta\eta)(i_0,\dots,i_k) \coloneqq d(\eta(i_0,\dots,i_k)) + (-1)^{|\eta|}\sum_{j=1}^{k-1}(-1)^j\eta(i_0,\dots,\widehat{i_j},\dots,i_k)$$

which lives in $\underline{\mathrm{Hom}}_k(E_{i_0},E_{i_k})$. We write $\delta = d + \Delta$. Here we define $|\eta| = 1$. The product is:

$$(\phi\circ\eta)(i_0,\dots,i_k) \coloneqq \sum_{j=0}^{k}(-1)^{|\phi|j}\phi(i_j,\dots,i_k)\circ\eta(i_0,\dots,i_j)$$

Both definitions follow section 5.2 of [10], with some corrections to the signs. We leave out the terms in $\delta\eta$ corresponding to leaving out i_0 and i_k as they do not live in the correct hom-spaces.
One can now check that $\Delta d = -d\Delta$ (and hence $\delta^2 = 0$) and we have the following Leibniz rule:

$$\delta(\phi\circ\eta) = (-1)^{|\eta|}(\delta\phi)\circ\eta + \phi\circ(\delta\eta)$$

The same equation holds for the summands d and Δ. (The unusual sign appears because of the backward notation for compositions.)

Example 1. For $n = 1$ we have $(\delta\eta + \eta^2)_{01} = d(\eta_{01}) + 0$, the expected cycle condition. For $n = 2$ we have for example

$$(\delta\eta + \eta^2)_{012} = d(\eta_{012}) + \eta_{02} - \eta_{12}\circ\eta_{01} \in \underline{\mathrm{Hom}}_1(E_0,E_2)$$

So an element of $\mathscr{D}_2$ is of the form (E,η) where $E = (E_0,E_1,E_2)$ and $\eta = (\eta_{01},\eta_{02},\eta_{12};\eta_{013})$ satisfies $d\eta + \eta^2 = 0$, which comes out to $d\eta_{ij} = 0$ and $d\eta_{012} = -\eta_{02} + \eta_{12}\circ\eta_{01}$. This agrees with our intuition that η_{012} is a homotopy from $\eta_{12}\circ\eta_{01}$ to η_{02}.

Morphisms from (E,η) to (F,ϕ) are given as follows.

$$\underline{\mathrm{Hom}}^{-m}_{\mathscr{C}_n}((E,\eta),(F,\phi)) = \{a(i_0,\dots,i_k)\}$$

where $a(i_0,\dots,i_k) \in \underline{\mathrm{Hom}}_{m-k}(E_{i_0},F_{i_k})$. We write $m = |a|$ for the degree of a morphism. We have a differential $d_{\eta,\phi}$ defined by

$$(d_{\eta,\phi}(a))(i_0,\dots,i_k) = \delta(a) + \phi\circ a - (-1)^{|a|}a\circ\eta$$

where composition and differential are defined as above. The Maurer–Cartan condition on η and ϕ together with the Leibniz rule ensures $(d_{\eta,\phi})^2 = 0$.

Example 2. For example $\mathscr{C}_1$ agrees with the path object in **dgCat** as constructed in Section 3 of [11]. Indeed, objects are homotopy invertible morphisms $\eta\colon A \to B$ and morphisms from η to ϕ are given by triples (a_0,a_1,a_{01}) with differential

$$\delta\colon (a_0,a_1,a_{01}) \mapsto (da_0, da_1, da_{01} + \phi\circ a_0 - (-1)^{|a_1|}a_1\circ\eta)$$

Note that there are induced face and degeneracy maps. The maps in the simplex category induce restriction functors $\partial_i \colon \mathscr{C}_n \to \mathscr{C}_{n-1}$ and inclusions $\sigma_i \colon \mathscr{C}_n \to \mathscr{C}_{n+1}$ that add an extra copy of E_i, connected by the identity map to E_i.
The replacement map $\iota : c\mathscr{C} \to \mathscr{C}_n$ is given by $(\sigma_0)^n$ in degree n.
Before we embark on the somewhat technical proof that $\mathscr{C}_\bullet$ is a simplicial resolution, we note the following application. We can extend the definitions of the differentials and composition to functions defined on general simplices. (That is, we replace "leaving out the i-th term" by the map induced by ∂_i etc.)

Proposition 6. *Given a simplicial set K we can construct $\mathscr{C}^K$ as the dg-category with objects (E, η) where $E \in (Ob\mathscr{C})^{K_0}$ and η assigns to every k-simplex in $K_{\geq 1}$ a map in $\underline{Hom}_{k-1}(E(\partial_0^k \sigma), E(\partial_{max}^k \sigma))$ satisfying the Maurer–Cartan equations. Hom-spaces are defined similarly to hom-spaces in $\mathscr{C}_\bullet$.*

Proof. This follows from the construction of $\mathscr{C}^K = \lim_{\Delta K^{\mathrm{op}}} \mathscr{C}_\bullet$. All the copies of $\mathscr{C}_n$ corresponding to degenerate simplices are themselves degenerate. □

Remark 6. Note that this shows that the construction of Morita cohomology in [4] as $K \mapsto \mathscr{C}^K$ corresponds to ∞-local systems as defined in [1].

Notation. Given an object or morphism α and a positive integer k we write $\alpha_{[k]}$ for the collection of all $\alpha_{i_0 \dots i_k}$.

Proposition 7. *The inclusion from the constant simplicial dg-category $c\mathscr{C}$ to $\mathscr{C}_\bullet$ is a levelwise weak equivalence.*

Proof. We have to check that the inclusion map $\iota \colon c\mathscr{C} \to \mathscr{C}_n$ is a quasi-equivalence.
Let us first show that ι induces weak equivalences on hom-complexes. We have to show that $\underline{\mathrm{Hom}}_{\mathscr{C}^{\Delta^n}}((\underline{E}, \eta), (\underline{F}, \phi)) \simeq \underline{\mathrm{Hom}}_{\mathscr{C}}(E, F)$ when both η and ϕ are of the form $(\mathbf{1}, 0)$, i.e. the constituent morphisms in degree 0 are the identity and all others are 0.
Write $(H, d_H) := \underline{\mathrm{Hom}}(E, F)$ and note that from the definitions we can write

$$\underline{\mathrm{Hom}}((E,0),(F,0)) \simeq (H[1] \otimes \bigwedge \langle e_0, \dots, e_n \rangle, D)$$

Here the e_i all have degree 1 and we identify $H.e_{i_0} \wedge \cdots \wedge e_{i_k}$ with the $a(i_0, \dots, i_k)$. The differential D is $d_H + \iota_{\sum e_i}$ where the second term denotes contraction. This complex is a resolution of (H, d_H).
Next we show ι is quasi-essentially surjective, i.e. show that any object (E, η) is equivalent to an object $(\underline{F}_0, (\mathbf{1}, 0))$ where $\underline{F}_0$ is of the form $(F_0, \dots, F_0)$.

We can deduce this if we can show that every (E, η) is equivalent to some (F, ϕ) such that all compositions which agree up to homotopy by $\delta\phi + \phi^2 = 0$ agree strictly, i.e. $\phi = (\phi_{[0]}, 0)$, and that any such $(F, (\phi_{[0]}, 0))$ is equivalent to $(\underline{F}_0, (\mathbf{1}, 0))$. The second part of this is immediate: We define a map from $(\underline{F}_0, (\mathbf{1}, 0))$ to $(F, (\phi_{[0]}, 0))$ by sending F_0 to F_i via $\phi(0, i) = \phi(i-1, i) \cdots \phi(0, 1)$. Since all $\phi(j, j+1)$ are homotopy invertible there is a homotopy inverse.
We will now show that any (E, η) is equivalent to (F, ϕ) where ϕ has no higher homotopies. Let $F = E$ and let $\phi(i, j) = \eta(j-1, j) \cdots \eta(i, i+1)$. We may assume by induction on n that all $\eta(i_0, \dots i_k)$ with $i_k < n$ are 0.
We define the homotopy equivalence $H \colon (E, \eta) \to (E, \phi)$ as follows:

$$
\begin{aligned}
H(i) &= \mathbf{1} \\
H(i_0, \dots, i_k) &= (-1)^{k-1} \eta(i_0, \dots, i_{k-1}, n-1, n) \quad \text{if } i_k = n \text{ and } i_{n-1} \neq n-1 \\
H(i_0, \dots, i_k) &= 0 \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{otherwise}
\end{aligned}
$$

And define H^- to be equal to H in degree 0 and $-H$ in degree > 0.
Then it is clear that H and H^- are inverses. Since $H(i_0, \dots, i_n)$ is zero unless $i_n = n$ there are no nontrivial compositions and the composition $\mathbf{1} \circ H(\dots)$ and $H^-(\dots) \circ \mathbf{1}$ cancel in degrees greater than 0.
So it remains to show that $dH = dH^- = 0$ to show we have a genuine homotopy equivalence.
We consider H first.
Putting together our definitions we find the following. Let us first assume $i_{k-1} \neq n-1$ and $i_k = n$. To obtain the correct signs recall that $|H| = 0$ and $|\eta| = |\phi| = 1$.

$$
\begin{aligned}
(dH)(i_0, \dots, i_k) = d(H(i_0, \dots, i_k)) &+ \sum_{j=1}^{k-1} (-1)^j H(i_0, \dots, \hat{i}_j, \dots, i_n) \\
&+ \sum_{j=0}^{k} (-1)^j \phi(i_j \dots i_n) \circ H(i_0, \dots, i_j) \\
&- \sum_{j=0}^{k} H(i_j, \dots, i_k) \circ \eta(i_0, \dots, i_j)
\end{aligned}
$$

This simplifies to:

$$\begin{aligned}(dH)(i_0,\dots,i_k) &= (-1)^{k-1}d\eta(i_0,\dots,n-1,n)\\ &\quad+(-1)^{k-2}\sum_j(-1)^j\eta(i_0,\dots,\hat{i}_j,\dots,n-1,n)\\ &\quad+0-(-1)^{k-2}\eta(i_1,\dots,n-1,n)\circ\eta(i_0,i_1)-\mathbf{1}\circ\eta(i_0,\dots,i_k)\\ &=0\end{aligned}$$

The last equality holds since the penultimate term is of the form

$$(-1)^{k-1}(\delta\eta+\eta^2)(i_0,\dots,i_{k-1},n-1,n)$$

This becomes clear if we write $\eta(i_0,\dots,i_k)=\eta(i_0,\dots,\widehat{n-1},n)$ and observe that all the other terms we expect in $\delta\eta+\eta^2$ are 0.
The other cases are easier. If $i_k\neq n$ all terms in the differential are 0 and if $i_{k-1}=n-1$ and $i_k=n$ there are only two nonzero terms, which cancel.
When we consider dH^- the sign of the term $\eta(i_0,\dots,i_k)$ changes, as it now comes from $\eta\circ H$ and not $H\circ\eta$. This cancels the effect of the sign of $H(i)$ also changing by a factor of -1. There are no other occurrences of the sign of $H(i)$ unless $k=1$ when all but the last two terms are zero and the last two terms cancel. □

Proposition 8. *The simplicial dg-category $\mathscr{C}_\bullet$ is Reedy fibrant.*

Proof. Write

$$\eta_{<n}\coloneqq(\eta_0,\dots,\widehat{\eta_{0\dots n}})=(\eta_{[0]},\dots,\eta_{[n-1]})$$

Then $M_n(\mathscr{C})$ is a subcategory of $\mathscr{C}_n$ whose objects are of the form $(E,\eta_{<n})$. In particular note that the Maurer–Cartan condition holds on all indexing sets except on $(0,\dots,n)$. Similarly, morphisms are of the form $s_{<n}$ where s is a morphism in $\mathscr{C}_n$. This is easily seen to be the correct limit, see Proposition 6. We write $\pi\colon\mathscr{C}_n\to M_n\mathscr{C}$ for the functor forgetting $\eta_{[n]}$.
It is immediate from the definition that there is a surjection on hom-spaces. So it remains to check the lifting property for homotopy invertible maps. We will first reduce to lifting contractions, as is done in the case of path objects in Section 3 of [11].
Note that by assumption the dg-category $\mathscr{C}$ is fibrant and hence has cones, cf. Section 2 of [11]. Then to see if a map h is homotopy invertible it suffices to check that $cone(h)$ is contractible.
So assume $h\colon(E,\eta_{<n})\to(F,\phi_{<n})$ is homotopy invertible in $M_n\mathscr{C}$ with homotopy inverse g and that $(E,\eta_{<n})$ is in the image of $\mathscr{C}_n$ under π. First we need to check that $(F,\phi_{<n})$ is also in the image of $\mathscr{C}_n$. It is enough to find $\phi_{[n]}$ such that $\delta\phi+\phi^2=0$ while we know that $\delta\phi_{<n}+\phi_{<n}^2=0$. In other words we are looking for $\phi_{[n]}$ such that $d\phi_{[n]}=(\Delta\phi+\phi^2)_{[n]}$.

We will first consider $g(n)\cdot(\Delta\phi+\phi^2)(0\dots n)$. Define $\rho\circ'\sigma$ to be $\rho\circ\sigma$ minus the term $\rho(n)\cdot\sigma(\cdots n)$. Then $-\circ'\sigma = -\circ\sigma$ if σ is η or ϕ. Note that d and Δ are compatible with $\circ'$ just as with the usual product.
Then $g(n)\cdot\phi(i\dots n) = (-g\circ'\phi+\eta\circ g-\delta g)(i\dots n)$ and we can perform the following computation, where we deduce the Maurer–Cartan condition in degree n from the Maurer–Cartan conditions in lower degrees.

$$\begin{aligned}
g(n)\cdot(\Delta\phi+\phi\circ\phi) &= -\Delta(g\circ'\phi)+\Delta(\eta\circ g)-\Delta\delta g\\
&\quad+(-g\circ'\phi+\eta\circ g-\delta g)\circ\phi\\
&= -g\circ'(\Delta\phi+\phi\circ\phi)+\eta\circ(\Delta g+g\circ\phi)\\
&\quad-dg\circ'\phi+\Delta\eta\circ g+d\Delta g\\
&\simeq g\circ' d\phi-dg\circ'\phi+\eta\circ\eta\circ g-\eta\circ dg+\Delta\eta\circ g\\
&= d(g\circ'\phi)-d\eta\circ g-\eta\circ dg\\
&\simeq -d(\eta\circ g)\\
&\simeq 0
\end{aligned}$$

Since $dh(n)=0$ we deduce that $h(n)g(n)(\Delta\phi+\phi^2)\simeq 0$ and it suffices to show $(h(n)g(n)-\mathbf{1})\cdot(\Delta\phi+\phi^2)\simeq 0$. We know there exists K with $dK = h(n)g(n)-\mathbf{1}$ so the desired homotopy follows if we can show that $d(\Delta\phi+\phi^2)=0$. One may check explicitly that $d(\Delta\phi) = -\Delta\phi\circ\phi+\phi\circ\Delta\phi$, using the fact that $d\phi = -\Delta\phi-\phi^2$ in degree less than n. Then we can use Maurer–Cartan in lower degrees again to deduce:

$$\begin{aligned}
d(\Delta\phi+\phi^2) &= d(\Delta\phi)-(-\Delta\phi-\phi^2)\circ\phi+\phi\circ(-\Delta\phi-\phi^2)\\
&= 0
\end{aligned}$$

Thus we know the domain and codomain of h are in the image of π and we can use surjectivity of hom-spaces to write $h=\pi(H)$. Now it suffices to show that the contraction of h lifts.
Let us assume we are given a contraction $s_{<n}$ of $cone(h) = (G,\gamma_{<n})$, we have to find a contraction s of (G,γ). By assumption we can write $d_\gamma(s_{<n}) = (\mathbf{1},0,\dots,0,t_{[n]})$ for some $t_{[n]}$. Now consider $0 = d_\gamma d_\gamma(s_{<n}) = (0,\dots,0,dt_{[n]}+0)$. This forces $\delta t_{[n]} = dt_{[n]} = 0$. But now we know that $ds_{[0]} = \mathbf{1}$ and hence $d\colon s_{[0]}t_{[n]}\mapsto t_{[n]}$ and $(s_{[0]},\dots,s_{[n-1]},s_{[0]}t_{[n]})$ is a contraction of (G,γ).
We deduce that H is contractible and the preimages of $(E,\eta_{<n})$ and $(F,\phi_{<n})$ are indeed homotopy equivalent. □

References

[1] J. Block and A. Smith, *A Riemann Hilbert correspondence for infinity-local systems*, ArXiv e-prints, (2009), 0908.2843v3.

[2] W. G. Dwyer and D. M. Kan, *Simplicial localizations of categories*, J. Pure Appl. Algebra, 17 (1980), pp. 267–284.
[3] P. S. Hirschhorn, *Model categories and their localizations*, vol. 99 of Mathematical Surveys and Monographs, American Mathematical Society, 2003.
[4] J. Holstein, *Morita Cohomology*. Preprint, 2014.
[5] K. Igusa, *Twisting cochains and higher torsion*, ArXiv e-prints, (2002), math/0212383.
[6] B. Keller, *On differential graded categories*, ArXiv e-prints, (2006), math/0601185.
[7] F. Muro, *Dwyer-Kan homotopy theory of enriched categories*, ArXiv e-prints, (2012), 1201.1575.
[8] G. Raptis, *On the Cofibrant Generation of Model Categories*, ArXiv e-prints, (2009), 0907.2726.
[9] C. Rezk, *Every homotopy theory of simplicial algebras admits a proper model*, Topology Appl., 119 (2002), pp. 65–94.
[10] C. Simpson, *Geometricity of the Hodge filtration on the ∞-stack of perfect complexes over X_{DR}*, ArXiv e-prints, (2005), math/0510269.
[11] G. Tabuada, *Homotopy theory of dg-categories via localizing pairs and Drinfeld's dg quotient*, Homology Homotopy Appl., 12 (2010), pp. 187–219.
[12] B. Toën, *Lectures on dg-categories*, in Topics in algebraic and topological K-theory, vol. 2008 of Lecture Notes in Math., Springer, Berlin, 2011, pp. 243–302.

Christ's College and University of Cambridge
E-mail address: J.V.S.Holstein@dpmms.cam.ac.uk